# Postpartum Depression and Risk Factors Among Females

Rabiya Fatima\* Manglesh Kumar\*\*

#### Abstract

**Background:** Postpartum depression (PPD) is a major depression with a postpartum onset. Various reasons, such as lack of support from the husband or the family, type of family, sex of the baby, past experiences, and many more, can lead to the onset of postpartum depression in a female. Aim: The aim of the present study was to assess postpartum depression and its possible risk factors among females.

Materials and Methods: The sample consisted of 100 postpartum females. The sample was selected by purposive sampling method from different nursing homes in Patna. Women with age ranging from 20 to 45 years were recruited as the sample of the study. After obtaining an informed consent, details of socio-demographic variables and risk factor related variables were gathered on interview with the postpartum females and postpartum depression was assessed using Edinburgh Postnatal Depression Scale.

**Result and Conclusion:** The result revealed that 58% of the females were suffering from postpartum depression. Possible risk factors identified in the study were mode of delivery, breast feeding problems, sleep disturbances, appetite disturbances, and mood swings during pregnancy. Age during first pregnancy was significantly negatively correlated with postpartum depression score.

Keywords: Postpartum depression, risk factors, female.

About Author: \*, Department of Psychological Sciences, Central University of South Bihar, Gaya \*\* Manglam, Department of Psychological Sciences, Central University of South Bihar, Gaya,

#### Introduction

Postpartum depression is an important public health concern, having a significant impact on the mother, the family, her partner, mother-infant interaction and on the long term emotional and cognitive development of the baby (Beck, 2006).

Postpartum Depression (PPD) can be defined as a depressive disorder which occurs within 4 weeks' postpartum, although a time frame of 3 months after birth also has been suggested for defining PPD (Elliott & Leverton, 2000). About one in every seven new mothers is affected by PPD (Wisner, 2006), resulting in an overall prevalence rate of 13% (O'Hara & Swain, 1996).

The prevalence of postpartum depression varies from 1.9% to 82.1% in developing countries and from 5.2% to 74.0%

in developed countries using self-reported questionnaire. The prevalence of postpartum depression was found to be 31.4% (Lakshmi et al., 2016). Prevalence of PPD was found to be 26.97%. Among the risk factors, advanced age, low socio-economic condition, minority community, fetal complications and vaginal delivery were found to be the most common (Khan and Basu, 2016).

Schmied et al. (2013) reported that mothers' past history of depression or existing mental disorders were associated with an increased risk of postpartum depression. Prior researchers have also found that lower maternal education level and poor family economic status were related to a higher prevalence of postpartum depression.

A cross sectional study was conducted by Kruthika et al.(2017) and reported that the prevalence of Post-natal depression was 41 (13.6%). Factors like age, literacy status, socio economic status, gravidity, sex of new born, mode of delivery and unplanned pregnancy were significantly associated with the prevalence of postnatal depression.

One hundred and eighty women were selected through consecutive sampling. The Edinburg Postnatal Depression Scale (EPDS) was administered to assess postnatal depression. Result showed that prevalence of probable depression was found to be 18%. Risk factors found to be significantly associated with postnatal depression were high risk pregnancy, being unhappy with in-laws, low income of the family, mood swings and low mood during pregnancy (Suguna et al., 2015).

A cross sectional study was done at Sri ManakulaVinayagar medical college and hospital, Pondicherry, India. Edinburgh Postnatal Depression Scale (EPDS) was used to detect postnatal depression. Type of marriage, recent stressful life event, addiction in husband, past history of psychiatric illness, order of pregnancy were significantly associated with PPD. In contrast to usual belief of parent's preference to male child, the study results showed most of them preferred female child (Poomalar and Bupathy, 2014).

The prevalence rate of postpartum depression among 200 Gujarati women was 12.5%. Women who delivered a girl child were observed to have higher chances of getting PPD. Women having past history of miscarriage, women feeling tense during pregnancy, women who could not confide in partner, and absence of someone other than mother and partner in whom the woman could confide were found to be the strongest predictors for developing postpartum depression (Desai et al., 2012). A survey revealed that one-third of women scoring within a depressive range at eight months postpartum were still depressed 12-18 months later, but only 15% sought help or were referred to a mental health professional (Lumley, 2005).In another study of 214 women, 86 reported high levels of depressive symptoms

(40.2%), but only 25 (11.7%) were actually diagnosed as being depressed (Hendrick, 2003).

In the Asian context, several important prevalence studies have been conducted. In Japan, of 70 Japanese mothers who were assessed by psychiatrists, Ueda et al. (2006) found that 27% mothers at 12 months postpartum were diagnosed as having experienced a new onset of depression. In Taiwan, Wang et al. (2003) conducted a study with 315 mothers six weeks post-delivery and found that about 31 and 12.9% mothers had experienced mild-to-moderate and moderateto-severe depression, respectively. In a recent study including 2,072 postpartum women in Malaysia, Yusuff et al. (2015) reported that 14.3% of mothers were categorized as having depression within the first 6 months postpartum. A study done in Tianjin, China indicated that the prevalence of postpartum depression was 10.2% among 463 mothers 35–60 days post-delivery (Zhang et al., 2001).

## Methods

**Aim:**The aim of the present was to assess postpartum depression and its possible risk factors among females

## **Objectives of the study:**

- To assess postpartum depression among females.
- To assess the possible risk factors of postpartum depression.

**Participants:** The sample consisted of 100 postpartum females. The sample was selected by purposive sampling method from different nursing homes in Patna. Women with age ranging from 20 to 45 years were recruited as the sample of the study. Women suffering from intellectual dysfunctions, women suffering from other psychiatric disorders, and women suffering from neurological disorders were excluded from the study. After obtaining an informed consent, details of socio-demographic variables and risk factor related variables were gathered on interview with the

postpartum females and postpartum depression was assessed using Edinburgh Postnatal Depression Scale.

#### **Tools:**

#### Socio-Demographic data sheet

A Socio-Demographic data sheetand risk factor related variables were especially designed for the present study like age, education, occupation, residence, income, order of pregnancy, miscarriage, past history of any mental illness or substance abuse, nature of delivery, term of pregnancy, sex of the child, support from husband and in-laws.

**Edinburgh Postnatal Depression Scale**(Cox et al., 1987): The 10-question Edinburgh Postnatal Depression Scale is a valuable and efficient way of identifying patients at risk for "perinatal" depression. The psychometric properties of the EPDS in primary health care were: 86% sensitivity, 78% specificity and 73% positive predictive value.

### **Procedure of Data Collection**

The researcher approached the concerned heads of the nursing homes and requested their permission to gain access to postpartum mothers. After obtaining an informed consent, details of socio-demographic variables and risk factor related variables were gathered on interview with the postpartum females and postpartum depression was assessed using Edinburgh Postnatal Depression Scale.

#### **Results-statistical analysis**

The collected data was analyzed by the statistical software Statistical Package for Social Sciences (SPSS) version 22.0 (for Windows). In descriptive statistics, mean and standard deviation were calculated for continuous variables while number and percentage were calculated for the discrete or categorical variables. To see the association among socio-demographic and clinical variables with depression, Pearson Correlation-Coefficient was used.

#### Results

## Table 1: Demographic Characteristics of the Sample (N= 100)

Variables	Mean ± SD
Age (in years)	$25.85 \pm 3.57$
Age during First Pregnancy (in years)	$22.82 \pm 3.21$

The mean age of the sample was found to be  $25.85 \pm 3.57$  years and mean age during first pregnancy was found to be  $22.82 \pm 3.21$  years.

Variables	n (%)	
	Below 10th	26 (26%)
Education	Above 10th	74 (74%)
	Urban	61 (61%)
Residence	Rural	39 (39%)

ISSN 0976 9218

	Nuclear	55 (55%)
Types of family	Joint	45 (45%)
	Employed	18 (18%)
Occupation	Unemployed	82 (82%)
	Satisfied	85 (85%)
Relationship with In-laws	Dissatisfied	15 (15%)
	Yes	6 (6%)
Substance addiction in husband	No	94 (94%)
	Male	61 (61%)
Sex of baby	Female	39 (39%)
	Yes	46 (46%)
Breast feeding problems	No	54 (54%)
	Yes	48 (48%)
Sleep disturbances	No	52 (52%)
	Yes	23 (23%)
Appetite disturbances	No	77 (77%)
	Yes	10 (10%)
Gender preference	No	90 (90%)
	Wanted	90 (90%)
Current pregnancy	Unwanted	10 (10%)
	Yes	13 (13%)
High risk pregnancy	No	87 (87%)
	Yes	94 (94%)
Mood swing during pregnancy	No	6 (6%)

Out of 100 sample, 26% had received education below 10th standard and 74% had received education above 10th standard, 61% belonged to urban areas and 39% belonged to rural areas, 55% belonged to nuclear families and 45% belonged to joint families, 18% were employed and 82% were unemployed, 15% had an unhealthy relationship with the in-laws, 6% reported substance addiction in husband, 61% had a male child and 39% had a female child, 46% had breast feeding problems, 48% had sleep disturbances, 23% had appetite disturbances, 10% had gender preference, 10% had unwanted pregnancy, 13% had high risk pregnancy, 94% had mood swings during

pregnancy, 69% had normal delivery and 31% had caesarean delivery, 9% had complications during delivery, and 8% reported having past history of miscarriage.

# Table 3:Clinical characteristics of females screened withpostpartum depression (N = 58)

Variables		n (%)
	Nuclear	34 (58.6%)
Type of Family	Joint	24 (41.4%)
	Male	36 (62.1%)
Sex of the Baby	Female	22 (37.9%)
	Normal	41 (70.7%)
Mode of Delivery	Caesarean	17 (29.3%)
Being Unhappy with	n-Laws	9 (15.5%)
Breast Feeding Proble	ems	29 (50%)
Sleep Disturbances		30 (51.7%)
Appetite Disturbances	\$	14 (24.1%)
Gender Preference		9 (15.5%)
Unwanted Pregnancy		9 (15.5%)
High Risk Pregnancy		12 (20.7%)
Mood Swings during	Pregnancy	55 (94.8%)

Out of 58 sample 34 (58. 6%) belonged to nuclear families and 24 (41.4%) belonged to joint families, 36 (62.1%) had a male child and 22 (37.9%) had a female child, 41 (70.7%) had normal delivery and 17 (29.3%) had caesarean delivery, 9 (15.5%) were unhappy with their in-laws, 29 (50%) had breast feeding problems, 30 (51.7%) had sleep disturbances, 14 (24.1%) had appetite disturbances, 9 (15.5%) had a gender preference, 9 (15.5%) had an unwanted pregnancy, 12 (20.7%) had high risk pregnancy and 55 (94.8%) had mood swings during pregnancy.

Fable 4: Distribution of sample according to cut off score of depression (13) and Mean and S	D
of EPDS Scores (N=100)	

EPDS Score	n (%)
Below 13	42 (42%)
Above 13	58 (58%)
	Mean ± SD
EPDS Total Score	$14.79 \pm 5.52$

ISSN 0976 9218

Forty two (42%) sample scored below 13, and 58 (58%) scored above 13. The mean of depression score was found to be  $14.79 \pm 5.52$ .

Table No.5: Correlation between EPDS score and Demographic Variable	les
---	-----

Variables	EPDS Score
Age	068
Age during First Pregnancy	366**

\*\* Correlation is significant at the <.01 level

Age during first pregnancy was significantly negatively correlated with postpartum depression score (r= -.366, p= <.01).

## Discussion

The present study indicated that 58% of the females had postpartum depression. Finding substantiates prior research done by Hendrick (2003), Lumley (2005), Lakshmi et al. (2016). Lumley (2005) conducted a survey which revealed that one-third of women scored within a depressive range. Research done by Lakshmi et al. (2016) revealed that 31.4% of the females were suffering with postpartum depression.

Various risk factors can lead to postpartum depression. The possible risk factors identified among the females screened with postpartum depression were type of family, marriage, sex of the baby, mode of delivery, being unhappy with in-laws, breast feeding problems, sleep disturbances, appetite disturbances, gender preference, unwanted pregnancy, high risk pregnancy, mood swings during pregnancy, complications during delivery. Almost similar findings were reported with other studies (Kruthika et al., 2017; Suguna et al., 2015).

In the present study 50% participants reported having breast feeding problems. The present finding is consistent with Rehman and Kazmi (2016). 51.7% reported having sleep disturbances and 24.1% had appetite disturbances. The present finding is supported by a prior research by Lakshmi et al. (2016) in which sleep disturbances, work and activity, changes in the appetite and general somatic

15.5% of the females reported having a gender preference. This finding is in line with the research by Poomalar and Bupathy (2014). In contrast to usual belief of parent's preference to male child, the study results showed most of them preferred female child. 20.7% of the females had a high-risk pregnancy while 94.8% reported having mood swings during pregnancy. The finding is substantiated by a prior research done by Suguna et al. (2015). Significant negative correlation was found between age during first pregnancy and

symptoms were observed to be more common

among the postpartum women with depression.

between age during first pregnancy and postpartum depression (r= -.366, p= <.01), indicating that first order of pregnancy at a young age can lead to postpartum depression. The experience of giving birth and caring for an infant can often be overwhelming for a young mother. First pregnancy at a young age often proves challenging to the mother because she might be immature and unskilled in caring and providing for a family. This may cause their perception towards their current situation as negative and overwhelming, and they may tend to be preoccupied with a sense of guilt and worthlessness. Hence, these feelings might lead to depression in young mothers.

## Conclusion

In the present study 58% of the females participants had postpartum depression. Significant negative correlation was found between age during first pregnancy and postpartum depression.Possible risk factors identified in the present study for postpartum depression were type of family, type of marriage, sex of the baby, mode of delivery, being unhappy with in-laws, breast feeding problems, sleeping disturbances, appetite disturbances, high risk pregnancy, mood swings during pregnancy. Limitations of the present study include sample size was small, additional psycho diagnostic tools can be administered for diagnostic confirmation of postpartum depression. Moreover, further studies are recommended to find out the factors leading to postpartum depression. There are a few future implications of the study.Early detection of and treatment for postpartum depression could benefit not only the mother's mental health and social functioning, but also the infant's physical and psychological health, and the family environment.Continuity of professional care from pregnancy to the postnatal period may also assist mothers who are reluctant to disclose that they may be experiencing postpartum depression.

## References

- Beck, C. T. (2006). Postpartum depression: It isn't just the blues. American Journal of Nursing, 106(5): 40-50.
- Cox, J. L., Holden, J. M., &Sagovsky, R. (1987). Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. British Journal of Psychiatry,150(6): 782-786.
- Desai, N., Mehta, R. Y., &Ganjiwale, J. (2012). Study of prevalence and risk factors of postpartum depression. National Journal of Medical Research, 2(2): 194-198.
- Elliott, S. A., &Leverton, T. J. (2000). Is the EPDS a magic wand? 'Myths' and the evidence base. Journal of Reproductive and Infant Psychology, 18(4): 297-307.
- Hendrick, V. (2003). Treatment of postnatal depression effective interventions are available, but the condition remains underdiagnosed. British Medical Journal, 327(7422): 1003-1014.
- Khan, B., &Basu, R. (2016). A study on evaluation of prevalence and risk factors of post-partum depression (PPD) in Indian

women. Journal of Dental and Medical Sciences, 15(7): 59-62.

- Kruthika, K., Udayar, S. E., &Mallapur, M. D. (2017). An epidemiological study of postnatal depression among women availing maternal health services in rural areas of Belagavi, Karnataka, India. International Journal of Community Medicine and Public Health. 4(3): 759-763.
- Lakshmi, B. G., Ramam, S., Devi, P. S., Raja Sanjay, B.B.P., S. R. K., & Rama Chandran S. (2016). Prevalence of postpartum depression at an Indian tertiary care teaching hospital. International Journal of Pharmaceutical and Clinical Research, 8(6): 616-618.
- Lumley, J. (2005). Attempts to prevent postnatal depression have not included mental health workers and have failed. British Medical Journal, 331(7507): 5-6.
- O'Hara, M.W., & Swain, A.M. (1996). Rates and risk of postpartum depression: A metaanalysis. International Review of Psychiatry, 8(1): 37–54.
- Poomalar, G. K., &Bupathy, A. (2014). Impact of socio-cultural factors on postpartum depression in South Indian women. International Journal of Reproduction, Contraception, Obstetrics and Gynaecology, 3(2): 338-343.
- Suguna, A., Naveen, R., &Surekha, A. (2015). Postnatal depression among women attending a rural maternity hospital in south India. National Journal of Community Medicine, 6(2): 297-301.
- Schmied, V., Johnson, M., Naidoo, N., Austin, M. P., Matthey, S., & Kemp, L., Mills, A., Meade, T., & Yeo, A. (2013). Maternal mental health in Australia and New Zealand: A review of longitudinal studies. Women Birth, 26(3): 167–178.
- Ueda, M., Yamashita, H., & Yoshida, K. (2006). Impact of infant health problems on post natal depression: Pilot study to evaluate a health visiting system. Psychiatry and Clinical Neuroscience, 60(2): 182–189.

Rabiya Fatima & Manglesh Kumar

- Wang, S. Y., Jiang, X. Y., Jan, W. C., & Chen, C. H. (2003). A comparative study of post natal depression and its predictors in Taiwan and China. American Journal of Obstetrics and Gynecology, 189(5): 1407–1412.
- Wisner, K. L., Chambers, C., & Sit, D. K. (2006). Postpartum depression: a major public health problem. Journal of American

Medical Association, 296(21): 2616–2618.

- Yusuff, A. S. M., Tang, L., Binns, C. W., & Lee, A. H. (2015). Prevalence and risk factors for post natal depression in Sabah, Malaysia: A cohort study. Women Birth, 28(1): 25–29.
- Zhang, X., Tu, Q., Xi, W., Jiang, Y., &Gao, Y. (2001). A study of incidence and the related factors of post-partum depression. China Journal of Psychiatry, 34: 236–238.